

NEONATAL

MICONAZOLE

This document should be read in conjunction with this **DISCLAIMER**

Unrestricted: Any prescriber may initiate treatment as per guideline

Presentation	Oral Gel: 2% (20mg/mL)		
Fieseillation	Cream: 2% (20mg/g)		
	Ointment: Miconazole 0.25% and Zinc oxide 15%		
	(combination product)		
Classification	Azole antifungal		
Indication	Treatment of topical and oral fungal infections		
Dose	Fungal infections		
	Oral Gel: Apply approximately 1cm of gel 4 times a day to the inside of the mouth. Rub around gums and inside of mouth using a cotton bud or finger. Ensure any excess gel is removed from the mouth. Note: Do not use a spoon to administer gel. Cream: Apply TWICE daily after nappy changes Ointment: Apply to entire affected area at each nappy change for at least 7 days.		
Guidelines & Resources	Breastfeeding challenges: Thrush in lactation		
Administration	Oral Gel: use Gel after feeds		
Adverse Reactions	Azole antifungals are generally well tolerated. Infrequently it may cause itch, burning, stinging or erythema.		
Storage	Store at room temperature - below 25°C		

Miconazole - Neonatal Page 1 of 2

References	Australian Medicines Handbook. Miconazole. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2019 [cited 2019 September]. Available from: https://amhonline.amh.net.au/
	MIMS Australia. Daktarin, Daktozin. In: MIMS Online [Internet]. St Leonards (New South Wales): MIMS Australia; 2019 [cited 2019 September]. Available from: https://www.mimsonline.com.au

Document owner:	Head of Department - Neonatology				
Author / Reviewer:	KEMH & PCH Pharmacy / Neonatology Directorate				
Date first issued:	October 2013	Version:	3.0		
Last reviewed:	October 2019	Next review date:	Oct 2022		
Endorsed by:	Neonatal Directorate Management Group	Date:	Oct 2019		
Standards Applicable:	NSQHS Standards: 1 Governance 3 Infection Control 4 Medication Safety;				
Printed or personally saved electronic copies of this document are considered uncontrolled.					
Access the current version from the WNHS website.					

© Department of Health Western Australia 2019