

NEONATAL

VITAMINS (Infant)

This document should be read in conjunction with this **DISCLAIMER**

Unrestricted: Any prescriber may initiate treatment as per guideline

| Presentation | Oral Solution: Penta-Vite® Multivitamin | | |
|----------------------|--|------------------|--|
| | Each 0.45mL contains: | | |
| | Vitamin A (retinol palmitate) | 390 micrograms | |
| | o Vitamin B1 (thiamine HCl) | 540 micrograms | |
| | o Vitamin B2 (riboflavine) | 810 micrograms | |
| | o Vitamin B (nicotinamide) | 7.1mg | |
| | o Vitamin B6 (pyridoxine) | 111.1 micrograms | |
| | Vitamin C (ascorbic acid) | 42.8 mg | |
| | Vitamin D3 (colecalciferol) | 10.1 microgram | |
| | Sodium saccharin | | |
| Classification | Multivitamin Supplement | | |
| Indication | Multivitamin supplement in < 35 weeks gestation on full feeds of unfortified breast milk | | |
| Dose | Oral: | | |
| | 0.45mL once a day | | |
| Administration | Oral: may be given at any time in regards to feeds | | |
| Adverse Reactions | Diarrhoea, nausea | | |
| Storage | Store unopened bottle at room temperature | | |
| _ | Once opened – refrigerate at 2-8°C. Discard 9 weeks after opening | | |
| References | <u>Vitamins and Mineral Supplements</u> | | |

Vitamins - Neonatal Page 1 of 2

| Document owner: | Head of Department - Neonatology | | | |
|---|--|-------------------|----------|--|
| Author / Reviewer: | KEMH & PCH Pharmacy / Neonatology Directorate | | | |
| Date first issued: | October 2001 | Version: | 3.0 | |
| Last reviewed: | May 2019 | Next review date: | May 2022 | |
| Endorsed by: | Neonatal Directorate Management Group | Date: | May 2019 | |
| Standards Applicable: | NSQHS Standards: 1 Governance 3 Infection Control 4 Medication Safety; | | | |
| Printed or personally saved electronic copies of this document are considered uncontrolled. | | | | |
| Access the current version from the WNHS website. | | | | |

© Department of Health Western Australia 2019

Vitamins - Neonatal Page 2 of 2